

LEXISCAN MYOCARDIAL PERFUSION IMAGING TEST

Name: _____

DOB: ____/____/____

Date of Test: ____/____/____

Time: ____:____ AM/PM

This test takes approximately 4 1/2 hours.

A Lexiscan Myocardial Perfusion Imaging test uses a radioactive imaging agent (not a dye) to produce pictures of the heart muscle. The imaging agent gives off a small amount of radiation which can be seen with a special camera. The amount of radiation exposure during this test is very small and poses no health risk. The test will help your doctor determine if there are areas of your heart which do not receive enough blood supply due to coronary artery disease. Areas of your heart which may have been damaged from a previous heart attack may also be seen. Prior to the test an IV will be started. During the rest portion of the test, you will receive an injection of the radioactive imaging agent, and pictures will be taken of your heart. For the stress portion of the test, a medication (Lexiscan) will be given through the IV to simulate exercise by dilating the blood vessels of the heart. During the infusion, you will be continuously monitored for blood pressure, heart rate, rhythm and ECG changes. You will be given an injection of the radioactive imaging agent during the infusion. Following the Lexiscan infusion, pictures will be taken to determine blood flow to the heart muscle.

Patient Instructions:

1. No caffeine for 24 hours prior to test. (This includes decaffeinated beverages, chocolate, and caffeine containing medications such as Anacin and Excedrin.)
2. Do not eat or drink anything other than water for two hours prior to the test.
No smoking for 6 hours prior to test.
3. Diabetics should eat a light breakfast and take their insulin.
4. If you have lung problems such as asthma, emphysema, or COPD, please let the nurse or technologist know prior to the test.
5. Take all your regular medications, unless otherwise instructed by your physician.
If you take any medications for breathing problems, please consult with the nurse or technologist. Any medications containing Aminophylline or Theophylline should not be taken for 48 hours prior to the test. If you take Persantine or Dipyridamole pills, please consult with the nurse or technologist.
6. Wear two-piece clothing and comfortable shoes. No shirts with metal snaps.
7. The ordering physician will be notified of the results. This physician will inform you of the results.

If you need to reschedule the test or have any questions regarding these instructions, please call our office.